

A re-evaluation of the lignocellulolytic *Agaricomycetes* from the Brazilian semi-arid region

E.R. DRECHSLER-SANTOS¹

T.B. GIBERTONI¹, A. GÓES-NETO² & M.A.Q. CAVALCANTI¹

drechslersantos@yahoo.com.br

¹Dept. Micologia, Universidade Federal de Pernambuco, Brazil

²Dept. Ciências Biológicas, Universidade Estadual de Feira de Santana, Brazil

Abstract — A checklist of the lignocellulolytic *Agaricomycetes* from the Brazilian semi-arid region is provided. It presents data on the distribution of 37 species, 10 families and six orders (*Agaricales*, *Auriculariales*, *Gloeophyllales*, *Hymenochaetales*, *Polyporales*, and *Russulales*). Twenty-eight taxa previously recorded from the studied region are excluded. The full checklist is available at www.mycotaxon.com/resources/weblists.html.

Key words: Caatinga, *Basidiomycetes*, *Aphyllophorales*

Introduction

Agaricomycetes (sensu Hibbett et al. 2007) or *Basidiomycetes* (sensu Kirk et al. 2001) comprises almost 21,000 species of *Basidiomycota* (Kirk et al. 2008) and includes wood-decomposing, parasitic, and ectomycorrhizal fungi (Hibbett 2006). The wood-decomposing fungi are termed lignocellulolytic because their enzymes are able to degrade wood hemi-cellulose, cellulose, and/or lignin, thus causing white and brown rots in plant species (Holf et al. 2004). Northeastern Brazil's semi-arid region (3–7° S and 35–45°W, 955.000 Km²) encompasses parts of nine States (FIG. 1). The Köppen classification of the climate is 'Bsh' (hot and dry), with an annual mean temperature 25.5°C and characterized by a short rainy (3–5 mos) and long dry (7–9 mos) season; the annual mean precipitation is 600 to 1045 mm (Moura & Ramos 2004). The area is dominated by Caatinga (seasonally dry tropical forest sensu Pennington et al. 2000), composed of typically tropophilous and thorny medium to low xerophytes, where succulent species of the plant families *Cactaceae*, *Euphorbiaceae*, and *Bromeliaceae* stand out in the landscape (Andrade-Lima 1981). Father Camille Torrend was the first researcher to collect and study lignocellulolytic *Agaricomycetes* in this Brazilian biome (Torrend 1940). The 18 *Agaricomycetes*




FIGURE 1. Brazilian semi-arid region (modified from: SPR/MI, www.mi.gov.br)

species taxonomically and nomenclaturally revised by Drechsler-Santos et al. (2008a) are deposited in URM, as cited by Maia & Gibertoni (2002). Other works that cover *Agaricomycetes* from this semi-arid region include Góes-Neto et al. (2003), Góes-Neto & Baseia (2006), and Drechsler-Santos et al. (2007). Considering the paucity of information on lignocellulolytic *Agaricomycetes* of Brazil's semi-arid region, this study represents a first comprehensive analysis of the diversity of lignocellulolytic *Agaricomycetes* in this unique biome.

Material and methods

This study was based on bibliographic research (Torrend 1940, Maia 1960, Góes-Neto 1999, Maia & Gibertoni 2002, Góes-Neto et al. 2003, Góes-Neto & Baseia 2006, Drechsler-Santos et al. 2007, 2008a) and revision of fungi exsiccata now curated in ALCB, CEPEC, HUEFS, and URM (Holmgren et al. 1990). The distribution of the species is provided according to occurrence in the States. Nomenclature and classification systems follow those of Hibbett et al. (2007), Index Fungorum (www.indexfungorum.org), and Centraalbureau voor Schimmelcultures (www.cbs.knaw.nl).

Results and discussion

The 37 lignocellulolytic *Agaricomycetes* species reported from Northeastern Brazil's Caatinga region represent 10 families and six orders. *Polyporales* is represented by 23 (62.2%) species, followed by eight (21.6%) species in the *Hymenochaetales*. The higher diversity of *Polyporales* agrees with results of other basidiomycete inventories in both Southern Brazilian subtropical (Drechsler-Santos et al. 2008b) and tropical Northeastern Brazilian Atlantic

forests (Gibertoni et al. 2004). *Polyporaceae* Fr. ex Corda is represented by the highest number of species (17, or 45.9% of the total), followed by 8 species in *Hymenochaetaceae* Imazeki & Toki. The genus with the highest number of species represented is *Phellinus* Quél. s.l. (six, 16.2%). Bahia is the State with highest number of species (33, 89.2%), of which 29 (78.4%) were recorded exclusively in the Bahian semi-arid. Pernambuco is represented by five (13.5%) species, of which *Ganoderma stipitatum* (Murrill) Murrill, *Gloeophyllum striatum* (Sw.) Murrill, and *Phylloporia chrysites* (Berk.) Ryvarden are reported only in this State. *Fomes fasciatus* (Sw.) Cooke was recorded only in Piauí. Only one species each is recorded for the Ceará and Paraíba semi-arid regions: *Favolus tenuiculus* P. Beauv. and *Pycnoporus sanguineus* (L.) Murrill, respectively. No representatives of lignocellulolytic *Agaricomycetes* have been reported from the semi-arid regions of Minas Gerais, Sergipe, Alagoas, or Rio Grande do Norte. Among the species identified thus far, only *Gloeophyllum striatum* causes brown rot; as observed in other tropical and subtropical areas (Nakasone 1996), all other (97.3%) species cause white-rot. We have excluded 28 taxa reported earlier because their exsiccata were not found or are in very poor condition (Drechsler-Santos et al. 2008a). The 37 species of this checklist represent our current knowledge of the diversity and distribution of lignocellulolytic *Agaricomycetes* in Northeastern Brazil's semi-arid region. Further investigations, however, will certainly increase the number fungal records and expand the reported species ranges throughout the area.

Acknowledgements

The authors would like to thank Mario Rajchenberg and Gregory Mueller for expert reviews of the manuscript, Leonor C. Maia, Maria A. Neves, and Domingos Cardoso for contribution to this work, and the herbarium curators of ALCB, CEPEC, HUEFS, and URM. This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq, Projects: Universal 478973/2006-3 and 479961/2007-7, Pesquisa em Biodiversidade do Semi-árido 010105.00/2004/PPBio, and Base de Dados Consolidada das Plantas e Fungos do Nordeste 552615/05-6. CNPq also provided a PhD scholarship to the first author and grants to A. Góes-Neto and M.A.Q. Cavalcanti.

Literature cited

- Andrade-Lima D. 1981. The Caatingas dominium. *Revista Brasileira de Botânica* 4: 149–163.
- Drechsler-Santos ER, Wartchow F, Baseia IG, Gibertoni TB, Cavalcanti MAQ. 2008a. Revision of the Herbarium URM I. *Agaricomycetes* from the semi-arid region of Brazil. *Mycotaxon* 104: 9–18.
- Drechsler-Santos ER, Groposo C, Loguerio-Leite C. 2008b. Additions to the knowledge of lignocellulolytic basidiomycetes in forests from Santa Catarina, Southern Brazil. *Mycotaxon* 103: 197–200 (in <http://mycotaxon.com/resources/weblists.html>).
- Drechsler-Santos ER, Vasconcellos-Neto JRT, Gibertoni TB, Góes-Neto A, Cavalcanti MAQ. 2007. Notes on *Navisporus*: *N. terrestris* and *N. floccosus* from Brazil. *Mycotaxon* 101: 265–269.

- Gibertoni TB, Ryvarden L, Cavalcanti MAQ. 2004. Poroid fungi (*Basidiomycota*) of the Atlantic rain forest in Northern Brazil. *Synopsis Fungorum* 18: 33–46.
- Góes-Neto A. 1999. Polypore diversity in the State of Bahia, Brazil: A historical review. *Mycotaxon* 72: 43–56.
- Góes-Neto A, Baseia IG. 2006. Filo *Basidiomycota*. In: *Diversidade e Caracterização dos Fungos do Semi-Árido Brasileiro* (eds.: LFP Gusmão, LC Maia). Associação Plantas do Nordeste II: 141–160.
- Góes-Neto A, Marques MFO, Andrade JD, Santos DS. 2003. Lignicolous aphyllorphoroid *Basidiomycota* in an Atlantic forest fragment in the semi-arid caatinga region of Brazil. *Mycotaxon* 88: 359–364.
- Hibbett DS. 2006. A phylogenetic overview of the *Agaricomycotina*. *Mycologia* 98(6): 917–925.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, Huhndorf S, James T, Kirk PM, Lücking R, Thorsten-Lumbsch H, Lutzoni F, Matheny PB, Mclaughlin DJ, Powell MJ, Redhead S, Schoch CL, Spatafora JW, Stalpers JA, Vilgalys R, Aime MC, Aptroot A, Bauer R, Begerow D, Benny GL, Castlebury LA, Crous PW, Dai YC, Gams W, Geiser DM, Griffith GW, Gueidan C, Hawksworth DL, Hestmark G, Hosaka K, Humber RA, Hyde KD, Ironside JE, Kõljalg U, Kurtzman CP, Larsson KH, Lichtwardt R, Longcore J, Miadlikowska J, Miller A, Moncalvo JM, Mozley-Standridge S, Oberwinkler F, Parmasto E, Reeb V, Rogers JD, Roux C, Ryvarden L, Sampaio JP, Schüßler A, Sugiyama J, Thorn RG, Tibell L, Untereiner WA, Walker C, Wang Z, Weir A, Weiss M, White MM, Winka K, Yao YJ, Zhang N. 2007. A higher-level phylogenetic classification of the *Fungi*. *Mycological Research* 111 (5): 509–547.
- Holf JA, Klopfenstein NB, Tonn JR, Mcdonald GI, Zambino PJ, Rogers JD, Peever TL, Carris LM. 2004. Roles of Woody Root-Associated Fungi in Forest Ecosystem Processes: Recent Advances in Fungal Identification. USDA Forest Service RMRS-RP-47, Rocky Mountain Research Station.
- Holmgren PK, Holmgren NH, Barnett LC. 1990. *Index herbariorum*. Part I: Herbaria of the World. 86^a ed. Bronx, New York Botanical Garden.
- Kirk PM, Cannon PF, David JC, Stalpers JA. 2001. *Dictionary of the Fungi*. 9^o Ed, CAB International.
- Kirk PM, Cannon PF, Minter DW, Stalpers Ja. 2008. *Ainsworth & Bisby's dictionary of the Fungi*. 10^o Ed, CAB International.
- Maia HS. 1960. Fungos Diversos. *Pulicações do Instituto de Micologia do Recife* 267: 1–51.
- Maia LC, Gibertoni TB. 2002. Fungos registrados no semi-árido nordestino. In: *Vegetação e Flora da Caatinga* (eds.: EVSB Sampaio, AM Giuliatti, J Virgínio, CFL Gamarra-Rojas). Recife, Associação Plantas do Nordeste/APNE e Centro Nordestino de Informações sobre Plantas/CNIP: 163–176.
- Moura AS, Ramos GCC. 2004. *Cenários para o Bioma Caatinga*. Conselho Nacional da Reserva da Biosfera da Caatinga, Secretaria de Ciência, Tecnologia e Meio Ambiente de Pernambuco, Recife.
- Nakasone KN. 1996. Diversity of lignicolous basidiomycetes in coarse woody debris. In: *Biodiversity and Coarse Woody Debris in Southern Forests: Effects on Biodiversity*. Proceedings of the Workshop (eds. J.W. McMim & D.A Crossley). USDA Forest Service: Southern Research Station: 35–42.
- Pennington RT, Prado DE, Pendry CA. 2000. Neotropical seasonally dry forests and Quaternary vegetation changes. *Journal of Biogeography* 27: 261–273.
- Torrend C. 1940. As Poliporáceas da Bahia e Estados Limítrofes. *Anais da 1ª Reunião Sul-Americana de Botânica* (1938) 2: 325–341.